


AD

TECHNICAL REPORT ARCCB-TR-95018

**IMPLEMENTING TENSOR ANALYSIS IN
Mathematica WITH ILLUSTRATIONS
FROM SCHWARZCHILD GRAVITATION**



L.V. MEISEL

MARCH 1995



**US ARMY ARMAMENT RESEARCH,
DEVELOPMENT AND ENGINEERING CENTER
CLOSE COMBAT ARMAMENTS CENTER
BENÉT LABORATORIES
WATERVLIET, N.Y. 12189-4050**



APPROVED FOR PUBLIC RELEASE; DISTRIBUTION UNLIMITED

19950710 050

DTIC QUALITY INSPECTED 5

DISCLAIMER

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

The use of trade name(s) and/or manufacturer(s) does not constitute an official indorsement or approval.

DESTRUCTION NOTICE

For classified documents, follow the procedures in DoD 5200.22-M, Industrial Security Manual, Section II-19 or DoD 5200.1-R, Information Security Program Regulation, Chapter IX.

For unclassified, limited documents, destroy by any method that will prevent disclosure of contents or reconstruction of the document.

For unclassified, unlimited documents, destroy when the report is no longer needed. Do not return it to the originator.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE March 1995		3. REPORT TYPE AND DATES COVERED Final	
4. TITLE AND SUBTITLE IMPLEMENTING TENSOR ANALYSIS IN <i>Mathematica</i> WITH ILLUSTRATIONS FROM SCHWARZCHILD GRAVITATION				5. FUNDING NUMBERS AMCMS: 611.02.H611.1	
6. AUTHOR(S) L.V. Meisel					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army ARDEC Benét Laboratories, AMSTA-AR-CCB-O Watervliet, NY 12189-4050				8. PERFORMING ORGANIZATION REPORT NUMBER ARCCB-TR-95018	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army ARDEC Close Combat Armaments Center Picatinny Arsenal, NJ 07806-5000				10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES Published in: <i>Computers in Physics</i>					
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited				12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Expressions that play roles in tensor analysis, such as Christoffel symbols and curvature tensors, are coded as <i>Mathematica</i> modules by straightforward transcription of their defining equations. The built-in functions can then be used to perform tensor analysis and numerical valuations. The utility of the <i>Mathematica</i> formulation is illustrated by examples from Schwarzschild gravitation.					
14. SUBJECT TERMS <i>Mathematica</i> , Tensor Analysis				15. NUMBER OF PAGES 13	
				16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UL		

TABLE OF CONTENTS

INTRODUCTION	1
PART I. TENSOR ANALYSIS IN <i>Mathematica</i>	1
Covariant Metric Tensor	1
Christoffel Symbols	2
The Curvature Tensor	3
The Ricci Tensor	4
PART II. APPLICATION: THE SCHWARZCHILD METRIC	6
Conventional Notation	6
The Einstein Equations	7
Solution of the Einstein Equations	7
The Schwarzschild Metric	9
REFERENCES	12

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	

INTRODUCTION.

Andrew and Fleming (ref 1) discussed the calculation of null geodesics for the Schwarzschild, Kerr-Newman, and Winicour metrics. Numerical computation employed FORTRAN code, which was produced by *Mathematica* (ref 2). The *Mathematica* code employed to generate the FORTRAN code for the neutral ($Q = 0$) Kerr-Newman geodesics was presented. Reference 1 provides a convincing demonstration of the utility of symbolic manipulation software for the development of error-free, complex FORTRAN (or C) code.

In this report, a general approach to problems in tensor analysis (ref 3) employing *Mathematica* is described. The standard expressions of tensor calculus are transcribed directly into *Mathematica* modules in Part I. The operation of the modules is illustrated as they are described by application to a simple two-dimensional curved space. The present formulation allows enumeration and simplification of complex tensor equations, employing built-in *Mathematica* functions (such as `Expand[]`, `Together[]`, or `Simplify[]`).

The code is applied to address some simple problems pertinent to Schwarzschild space-time in Part II. Employing the built-in *Mathematica* operator `Simplify[]`, the modules described produced FORTRAN coded Runge-Kutta geodesic equations for Kerr-Newman space-time in less than thirty minutes on a 386-based PC operating at 30 MHz. Of course, if simplification based on more efficient *Mathematica* operations (e.g., combinations of `Apart[]`, `Expand[]`, and `Together[]`) can be achieved, symbolic computation times can be dramatically reduced. Computation times are much shorter for the Schwarzschild geodesic equations.

PART I. TENSOR ANALYSIS IN *Mathematica*

Covariant Metric Tensor

The starting point for a typical problem in tensor analysis is the specification of the (covariant) metric tensor.

The operation of the *Mathematica* tensor analysis modules defined here is demonstrated by applications to geometry on the surface of a helicoid in R^3 having

$$g = \|g_{ij}\| = \begin{pmatrix} 1 & 0 \\ 0 & c^2 + x[1][s]^2 \end{pmatrix}$$

where the coordinates are $x[1][s]$ and $x[2][s]$ and c is constant. This g is referred to as the "example metric" in Part I. One encodes this g directly into *Mathematica* via

$$g = \text{DiagonalMatrix}[\{1, c^2 + x[1][s]^2\}];$$

Note that coordinates are given in the form $x[i][s]$. For the current analysis, one could simply use $x[i]$. However, as in Part II, one is often interested in derivatives with respect to the line element s , and we choose to include this dependence from the beginning.

Christoffel Symbols

The first step in a tensor problem is the computation of the Christoffel symbols. Christoffel symbols of the first kind are defined as

$$\Gamma_{i\alpha\beta} = \Gamma_{i\beta\alpha} = \frac{1}{2} \left(\frac{\partial g_{ai}}{\partial x^\beta} + \frac{\partial g_{i\beta}}{\partial x^\alpha} - \frac{\partial g_{\alpha\beta}}{\partial x^i} \right)$$

Christoffel symbols of the second kind are given by

$$\Gamma_{\alpha\beta}^\delta = \sum_i g^{\delta i} \Gamma_{i\alpha\beta} = g^{\delta i} \Gamma_{i\alpha\beta}$$

where the last equality introduces "summation convention" which is adopted in this report and the contravariant metric tensor g^{ij} is the inverse of the covariant metric, i.e.,

$$\delta_\beta^\alpha = g^{\alpha i} g_{i\beta}$$

The following *Mathematica* module defines the contravariant metric tensor and Christoffel symbols for arbitrary metric tensors in spaces of arbitrary dimension.

```

setup[g_]:=Module[{dim=Length[g]},
  ginv =Together[Inverse[g]];
  GAMMA =Array[g1,{dim,dim,dim}];
  gamma =Array[g2,{dim,dim,dim}];
  Do[ Do[ Do[GAMMA[[i,k,j]]=GAMMA[[i,j,k]]=
    Together[(1/2)(D[g[[i,j]],x[k][s]]+
      D[g[[k,i]],x[j][s]]-D[g[[j,k]],x[i][s]])),
    {i,dim},{k,j,dim},{j,dim}];
  Do[ Do[ Do[gamma[[i,k,j]]=gamma[[i,j,k]]=
    Together[Sum[ginv[[i,l]]GAMMA[[l,j,k]],{l,dim}]],
    {i,dim},{k,j,dim},{j,dim}]] ]

```

N. B., the partial derivative of a function $f[u,v,\dots,x,\dots]$ with respect to the variable x is computed via $D[f[u,v,\dots,x,\dots],x]$ in *Mathematica*. N.B., the *Mathematica* `Together[]` operator was effective in simplifying the resulting expressions for a number of "simple problems." Of course, the user could apply an other built-in (e.g., `Simplify[]`) or user-defined *Mathematica* operator.

After `setup[g]` is run, the contravariant metric tensor is returned as `ginv`, Christoffel symbols of the first kind are in the array `GAMMA`, i.e., $\Gamma_{ijk} = \text{GAMMA}[[i,j,k]]$, and Christoffel symbols of the second kind are in the array `gamma`, i.e., $\Gamma_{jk}^i = \text{gamma}[[i,j,k]]$.

To compute the Christoffel symbols and g^{-1} for the test metric, enter

setup[g].

Example: Display gamma for the example metric,

gamma

returns

$$\{\{0, 0\}, \{0, -\frac{x[1][s]}{c + x[1][s]}\}, \{\{0, \frac{x[1][s]}{c + x[1][s]}\}, \{0\}\}, \{0\}\},$$

$$\{\{0, 0\}, \{0, -\frac{x[1][s]}{c + x[1][s]}\}\},$$

Exercise: Code a *Mathematica* module to compute covariant derivatives.

The Curvature Tensor

The curvature tensor can be computed directly from the $\Gamma_{\alpha\beta}^i$

$$B_{\alpha j k}^i = \frac{\partial \Gamma_{\alpha j}^i}{\partial x^k} - \frac{\partial \Gamma_{\alpha k}^i}{\partial x^j} + \Gamma_{\alpha j}^{\beta} \Gamma_{\beta k}^i - \Gamma_{\alpha k}^{\beta} \Gamma_{\beta j}^i$$

Transcription of the expressions for the $B_{\alpha j k}^i$ is straightforward,

```
curvature[gamma_,i_,a_,j_,k_] := Module[{b,dim=Length[gamma]},
  Together[
    D[gamma[[i,a,j]],x[k][s]]-D[gamma[[i,a,k]],x[j][s]]+
    Sum[gamma[[b,a,j]]gamma[[i,b,k]]-
      gamma[[b,a,k]]gamma[[i,b,j]],{b,dim} ] ]
```

and full curvature tensor is returned by

```
curvature[gamma_] := Module[{dim=Length[gamma]},
  B=Array[r1,{dim,dim,dim,dim}];
  Do[Do[Do[Do[B[[i,a,j,k]]=curvature[gamma,i,a,j,k],
    {i,dim}],{a,dim}],{j,dim}],{k,dim}];B]
```

The covariant form, $R_{hijk} = g_{ha} B_{i j k}^a$ is called the Riemann-Christoffel curvature tensor.

Example: Display the curvature tensor for the example metric

B = curvature[gamma]

returns

$$\begin{aligned} & \left\{ \left\{ \left\{ 0, 0 \right\}, \left\{ 0, 0 \right\} \right\}, \left\{ \left\{ 0, -\frac{c^2}{c^2 + x[1][s]^2} \right\}, \left\{ -\frac{c^2}{c^2 + x[1][s]^2}, 0 \right\} \right\} \right\}, \\ & \left\{ \left\{ 0, -\frac{c^2}{c^2 + x[1][s]^2} \right\}, \left\{ -\frac{c^2}{c^2 + x[1][s]^2}, 0 \right\} \right\}, \left\{ \left\{ 0, 0 \right\}, \left\{ 0, 0 \right\} \right\} \right\} \end{aligned}$$

The Ricci Tensor

The Ricci tensor plays an important role in relativity theory. It is defined as a contraction of the curvature tensor,

$$R_{ij} = B_{iaj}^a = \frac{\partial \Gamma_{ia}^\alpha}{\partial x^j} - \frac{\partial \Gamma_{ij}^\alpha}{\partial x^a} + \Gamma_{ia}^\beta \Gamma_{\beta j}^\alpha - \Gamma_{ij}^\beta \Gamma_{\beta a}^\alpha$$

Thus, to Print[] the components of $R_{ij} = B_{iaj}^a$, enter the *Mathematica* statement
 Do[Do[Print["R(",i,"",j,"") = ",Simplify[Sum[B[[a,i,a,j]],{a,2}]],
 {i,2}], {j,2}]

which yields

$$\begin{aligned} R(1,1) &= \frac{c^2}{(c^2 + x[1][s]^2)^2} \\ R(2,1) &= 0 \\ R(1,2) &= 0 \\ R(2,2) &= \frac{c^2}{c^2 + x[1][s]^2} \end{aligned}$$

N.B., Sum[*expression*,{i,n}] returns the sum of *expression* evaluated for i-values running from 1 to (the integer) n. Do[*expression*,{i,n}] works similarly.

We also transcribe separate *Mathematica* modules to compute the Ricci tensor,

```
Ricci[gamma_,i_,j_]:=Module[{m,n,dim=Length[gamma]},
Together[Sum[D[gamma[[m,i,m]],x[j][s]]-D[gamma[[m,i,j]],x[m][s]]+
Sum[gamma[[n,i,m]]gamma[[m,n,j]]-
gamma[[n,i,j]]gamma[[m,n,m]],{n,dim}},{m,dim}]]]
```

and

```
Ricci[gamma_]:=Module[{dim=Length[gamma]},
R=Array[r1,{dim,dim}];
Do[Do[R[[i,j]]=Ricci[gamma,i,j],{i,dim}},{j,dim}];R]
```

Example: Direct computation of Ricci tensor for the example metric.

```
Ricci[gamma]
```

returns

$$\left\{ \left\{ \frac{c^2}{(c^2 + x[1][s]^2)}, 0 \right\}, \left\{ 0, \frac{c^2}{c^2 + x[1][s]^2} \right\} \right\}$$

which is identical to the results obtained by contraction of B, as expected.

Exercise: Compute the Christoffel symbols and Ricci tensor for

$$\|g_{ij}\| = \begin{vmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{vmatrix}$$

PART II. APPLICATION: THE SCHWARZCHILD METRIC

Development of a spherically-symmetric $\|g_{ij}\|$ consistent with the Einstein equations.
Assumption: $\|g_{ij}\|$ is spherically-symmetric. Assume that

$$g = \|g_{ij}\| = \begin{pmatrix} L[r] & 0 & 0 & 0 \\ 0 & -r^2 & 0 & 0 \\ 0 & 0 & -r^2 \sin^2(\theta) & 0 \\ 0 & 0 & 0 & M[r] \end{pmatrix}$$

$$= \begin{pmatrix} L[x[1][s]] & 0 & 0 & 0 \\ 0 & -x[1][s]^2 & 0 & 0 \\ 0 & 0 & -x[1][s]^2 \sin^2(x[2][s]) & 0 \\ 0 & 0 & 0 & M[x[1][s]] \end{pmatrix}$$

where the functions $L[r]$ and $M[r]$ are to be determined. Encode via,

`g=DiagonalMatrix[{L[x[1][s]],-x[1][s]^2,-(x[1][s]Sin[x[2][s]])^2,M[x[1][s]]}];`

Conventional Notation

Direct substitution is achieved in *Mathematica*, via the construction

`expression/.u->v`

which returns *expression* with occurrences of *u* replaced by *v*. Thus, to present output in {r,q,j,t}-form, define the following replacement rules:

`rt={x[1][s]->r,x[2][s]->q,x[3][s]->j,x[4][s]->t};`

N.B., *Mathematica* 2.1 does *not* have Greek fonts. We substitute *j* for ϕ and *q* for θ here and in the sequel with a text editor.

Compute the Christoffel symbols, etc.

`setup[g];`

The Einstein Equations

The Einstein equations require that $R_{ij} = 0$. Compute and display the Ricci tensor.

$$R = \text{Simplify}[\text{Ricci}[\text{gamma}]]/.rt$$

returns the Ricci tensor for the spherically-symmetric metric function,

$$\begin{aligned} & \left\{ \left\{ -\frac{L'[r]}{r L[r]} - \frac{L'[r] M'[r]}{4 L[r] M[r]} - \frac{M'[r]^2}{4 M[r]^2} + \frac{M''[r]}{2 M[r]}, 0, 0, 0 \right\}, \right. \\ & \left\{ 0, -1 - \frac{1}{L[r]} + \frac{r L'[r]}{2 L[r]^2} + \frac{r M'[r]}{2 L[r] M[r]}, 0, 0 \right\}, \\ & \left\{ 0, 0, (\sin[T] (-2 L[r] M[r] - 2 L[r]^2 M[r] + \right. \\ & \left. r M[r] L'[r] - r L[r] M'[r])) / (2 L[r]^2 M[r]), 0 \right\}, \\ & \left. \left\{ 0, 0, 0, \frac{M'[r]}{r L[r]} - \frac{L'[r] M'[r]}{4 L[r] M[r]} - \frac{M'[r]^2}{4 L[r] M[r]^2} + \frac{M''[r]}{2 L[r]} \right\} \right\} \end{aligned}$$

Solution of the Einstein Equations

We seek functions $L[r]$ and $M[r]$ such that the Einstein equations are satisfied.

Step 1. Eliminate the $M''[r]$ terms from the 1,1 and 4,4 parts of $R_{ij} = R[[i,j]] = 0$.

$$\text{temp} = ((\text{Simplify}[(R[[1,1]]M[r[s]] - R[[4,4]]L[r[s]])/.rt]) == 0)$$

returns

$$-\left(\frac{M[r] L'[r] + L[r] M'[r]}{r L[r]} \right) == 0$$

Step 2. Solve for $L[r]$ in terms of $M[r]$. Use

$$\|g_{ij}\| \xrightarrow{r \rightarrow \infty} \begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

and the *Mathematica* operator `DSolve[]` to obtain Rule0:

```
Rule0=ExpandAll[Flatten[DSolve[
  {L[Infinity]==-1,M[Infinity]==1,temp},L[r],r]]]
```

Mathematica issues a "warning:"

Solve::ifun:

Warning: Inverse functions are being used by Solve, so some solutions may not be found.

then returns the solution of interest

$$\{L[r] \rightarrow -\frac{1}{M[r]}\}$$

Step 3. *Mathematica* will not apply the rule for $L[r]$ to replace $L'[r]$. Thus, one specifies a separate rule for $L'[r]$ and defines Rule1 as follows:

```
Rule1=Flatten[{Rule0,Solve[temp,L'[r]]/.Rule0}]/.r->x[1][s]
```

which returns

$$\{L[x[1][s]] \rightarrow -\frac{1}{M[x[1][s]]}, L'[x[1][s]] \rightarrow \frac{M'[x[1][s]]}{M[x[1][s]]}\}$$

N.B., `Flatten[{{u},{v}}]` returns $\{u,v\}$, etc. Thus, the subrules comprising Rule1 will be applied consecutively.

Step 4. Use Rule1 and $R_{22} = 0$ to define the form of M . Express M in a rule (Rule2).

```
Rule2=Flatten[DSolve[
  0==Factor[Expand[R[[2,2]]/.Rule1]]/.rt,M,r]]]
```

which returns the rule governing the function M ,

$$\{M \rightarrow (1 + \frac{C[1]}{x[1][s]^2})\}$$

where C[1] is a constant of integration. The rule on the function works as expected, i.e.,

returns

$$1 + \frac{C[1]}{x[1][s]^2}$$

The Schwarzschild Metric

Applying Rule1 and Rule2, one obtains the form of the spherically-symmetric metric tensor consistent with $R = 0$:

returns

$$\left\{ \left(-\frac{C[1]}{1 + \frac{C[1]}{r^2}} \right), 0, 0, 0 \right\}, \left\{ 0, -r^2, 0, 0 \right\}, \left\{ 0, 0, 0, 1 + \frac{C[1]}{r^2} \right\}$$

As expected, this is the Schwarzschild metric form. The usual form of the metric tensor is obtained by identifying the integration constant C[1] with $-2 G M/c^2$, where G is the gravitational constant, M is the central mass, and c is the speed of light.

Exercise: Demonstrate that $R = 0$ for Schwarzschild gravitation.

Exercise: The geodesic trajectories are given by solutions of the geodesic equations

$$\frac{d^2 x[i][s]}{ds^2} + \Gamma_{\alpha\beta}^i \frac{dx[\alpha][s]}{ds} \frac{dx[\beta][s]}{ds} = 0$$

Derive the geodesic equations for

$$\|g_{ij}\| = \begin{vmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{vmatrix}$$

Exercise: Derive the geodesic equations for the "example metric,"

$$\|g_{ij}\| = \begin{vmatrix} 1 & 0 \\ 0 & x[1][s]^2 + c^2 \end{vmatrix}$$

Exercise: Planetary orbits.

1. Derive the geodesic equations for the Schwarzschild metric.
2. Demonstrate that if $q[0] = p/2$ and $q'[0] = 0$, then $q[s] = p/2$.
 - a. Express the geodesic equations for $q[s] = p/2$.
3. Show that $r[s]^2 j'[s] = h = \text{constant}$.
4. Show that $(1 - (2G M/c^2) / r[s]) t'[s] = K = \text{constant}$.
5. Use the results of parts 1 through 4 and

$$1 = \frac{ds^2}{ds^2} = g_{ij} \frac{dx[i][s]}{ds} \frac{dx[j][s]}{ds}$$

$$= g_{ij} x[i]'[s] x[j]'[s] \rightarrow \sum_{i=1}^4 g_{ij} x[i]'[s]^2$$

to derive the *Mathematica* expression

$$1 == \frac{\frac{2}{K} - \frac{2}{h} (x[1])'[s]^2}{1 + \frac{C[1]}{x[1][s]} - \frac{2}{x[1][s]} + \frac{C[1]}{x[1][s]}}$$

6. Make the change of variables,

$$x[1][s] \rightarrow 1/u[x[3][s]],$$

and employ the condition derived in 3, to establish that

$$(x[1])'[s] \rightarrow -(h u'[x[3][s]])$$

and derive the equations:

$$1 == -(h^2 u[\phi]^2) + \frac{K^2}{1 + C[1] u[\phi]} - \frac{h^2 u'[\phi]^2}{1 + C[1] u[\phi]}$$

$$u'[\phi]^2 + u[\phi]^2 == -h^2 + \frac{-2 K C[1] u[\phi]}{h^2} - C[1] u[\phi]^3$$

and

$$u''[\phi] + u''[\phi] == -\left(\frac{C[1]}{2}\right) - \frac{3 C[1] u[\phi]^2}{2}$$

The last equation for $u[\phi]$ ($= 1/r[j]$) frequently serves as the basis for a discussion of the advance of the perihelion of planetary orbits. See, for example, Crandall (ref 4). How should this approach be modified to treat photon trajectories?

Exercise: Compute the Christoffel symbols, curvature tensor, Ricci tensor, and geodesic equations for charge-free (i.e., $Q = 0$) Kerr-Newman gravitation. Demonstrate that $R = 0$ for the Kerr-Newman metric.

REFERENCES

1. Keith Andrew and Charles G. Fleming, *Comp. in Phys*, Vol. 6, 1992, p. 498.
2. Stephen Wolfram, *Mathematica, A System for Doing Mathematics by Computer*, Addison-Wesley, Redwood City, CA, 1991.
3. The expressions presented in this report may be found in any textbook that addresses Riemann geometry. See, for example, Harry Lass, *Vector and Tensor Analysis*, The Maple Press, York, PA, 1950.
4. Richard E. Crandall, *Mathematica for the Sciences*, Addison-Wesley, Redwood City, CA, 1991.

TECHNICAL REPORT INTERNAL DISTRIBUTION LIST

	<u>NO. OF COPIES</u>
CHIEF, DEVELOPMENT ENGINEERING DIVISION	
ATTN: AMSTA-AR-CCB-DA	1
-DB	1
-DC	1
-DD	1
-DE	1
CHIEF, ENGINEERING DIVISION	
ATTN: AMSTA-AR-CCB-E	1
-EA	1
-EB	1
-EC	
CHIEF, TECHNOLOGY DIVISION	
ATTN: AMSTA-AR-CCB-T	2
-TA	1
-TB	1
-TC	1
TECHNICAL LIBRARY	
ATTN: AMSTA-AR-CCB-O	5
TECHNICAL PUBLICATIONS & EDITING SECTION	
ATTN: AMSTA-AR-CCB-O	3
OPERATIONS DIRECTORATE	
ATTN: SMCWV-ODP-P	1
DIRECTOR, PROCUREMENT & CONTRACTING DIRECTORATE	
ATTN: SMCWV-PP	1
DIRECTOR, PRODUCT ASSURANCE & TEST DIRECTORATE	
ATTN: SMCWV-QA	1

NOTE: PLEASE NOTIFY DIRECTOR, BENÉT LABORATORIES, ATTN: AMSTA-AR-CCB-O OF ADDRESS CHANGES.

TECHNICAL REPORT EXTERNAL DISTRIBUTION LIST

	<u>NO. OF COPIES</u>		<u>NO. OF COPIES</u>
ASST SEC OF THE ARMY RESEARCH AND DEVELOPMENT ATTN: DEPT FOR SCI AND TECH THE PENTAGON WASHINGTON, D.C. 20310-0103	1	COMMANDER ROCK ISLAND ARSENAL ATTN: SMCRI-ENM ROCK ISLAND, IL 61299-5000	1
ADMINISTRATOR DEFENSE TECHNICAL INFO CENTER ATTN: DTIC-OC (ACQUISITION GROUP) BLDG. 5, CAMERON STATION ALEXANDRIA, VA 22304-6145	2	MIAC/CINDAS PURDUE UNIVERSITY P.O. BOX 2634 WEST LAFAYETTE, IN 47906	1
COMMANDER U.S. ARMY ARDEC ATTN: SMCAR-AEE	1	COMMANDER U.S. ARMY TANK-AUTMV R&D COMMAND ATTN: AMSTA-DDL (TECH LIBRARY) WARREN, MI 48397-5000	1
SMCAR-AES, BLDG. 321	1	COMMANDER U.S. MILITARY ACADEMY ATTN: DEPARTMENT OF MECHANICS	1
SMCAR-AET-O, BLDG. 351N	1	WEST POINT, NY 10966-1792	
SMCAR-FSA	1	U.S. ARMY MISSILE COMMAND REDSTONE SCIENTIFIC INFO CENTER	2
SMCAR-FSM-E	1	ATTN: DOCUMENTS SECTION, BLDG. 4484 REDSTONE ARSENAL, AL 35898-5241	
SMCAR-FSS-D, BLDG. 94	1	COMMANDER U.S. ARMY FOREIGN SCI & TECH CENTER ATTN: DRXST-SD	1
SMCAR-IMI-I, (STINFO) BLDG. 59	2	220 7TH STREET, N.E. CHARLOTTESVILLE, VA 22901	
PICATINNY ARSENAL, NJ 07806-5000		COMMANDER U.S. ARMY LABCOM MATERIALS TECHNOLOGY LABORATORY ATTN: SLCMT-IML (TECH LIBRARY)	2
DIRECTOR U.S. ARMY RESEARCH LABORATORY ATTN: AMSRL-DD-T, BLDG. 305	1	WATERTOWN, MA 02172-0001	
ABERDEEN PROVING GROUND, MD 21005-5066		COMMANDER U.S. ARMY LABCOM, ISA ATTN: SLCIS-IM-TL	1
DIRECTOR U.S. ARMY RESEARCH LABORATORY ATTN: AMSRL-WT-PD (DR. B. BURNS)	1	2800 POWER MILL ROAD ADELPHI, MD 20783-1145	
ABERDEEN PROVING GROUND, MD 21005-5066			
DIRECTOR U.S. MATERIEL SYSTEMS ANALYSIS ACTV ATTN: AMXSY-MP	1		
ABERDEEN PROVING GROUND, MD 21005-5071			

NOTE: PLEASE NOTIFY COMMANDER, ARMAMENT RESEARCH, DEVELOPMENT, AND ENGINEERING CENTER, BENÉT LABORATORIES, CCAC, U.S. ARMY TANK-AUTOMOTIVE AND ARMAMENTS COMMAND, AMSTA-AR-CCB-O, WATERVLIET, NY 12189-4050 OF ADDRESS CHANGES.

TECHNICAL REPORT EXTERNAL DISTRIBUTION LIST (CONT'D)

	<u>NO. OF COPIES</u>		<u>NO. OF COPIES</u>
COMMANDER		WRIGHT LABORATORY	
U.S. ARMY RESEARCH OFFICE		ARMAMENT DIRECTORATE	
ATTN: CHIEF, IPO	1	ATTN: WL/MNM	1
P.O. BOX 12211		EGLIN AFB, FL 32542-6810	
RESEARCH TRIANGLE PARK, NC 27709-2211			
DIRECTOR		WRIGHT LABORATORY	
U.S. NAVAL RESEARCH LABORATORY		ARMAMENT DIRECTORATE	
ATTN: MATERIALS SCI & TECH DIV	1	ATTN: WL/MNMF	1
CODE 26-27 (DOC LIBRARY)	1	EGLIN AFB, FL 32542-6810	
WASHINGTON, D.C. 20375			

NOTE: PLEASE NOTIFY COMMANDER, ARMAMENT RESEARCH, DEVELOPMENT, AND ENGINEERING CENTER,
BENET LABORATORIES, CCAC, U.S. ARMY TANK-AUTOMOTIVE AND ARMAMENTS COMMAND,
AMSTA-AR-CCB-O, WATERVLIET, NY 12189-4050 OF ADDRESS CHANGES.
